

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/353889466>

Association between Cluster of Differentiation 24 (CD24) Polymorphism, Talin-1 gene expression and Hepatocellular Carcinoma prevalence in Egyptian population

Article in Archives of Medical Science · August 2021

DOI: 10.5114/aoms/140451

CITATIONS

0

READS

8

4 authors, including:



Mohamed M Hafez

Ahram Canadian University

5 PUBLICATIONS 11 CITATIONS

[SEE PROFILE](#)



Lina Jamil

October 6 University

14 PUBLICATIONS 50 CITATIONS

[SEE PROFILE](#)

Association between Cluster of Differentiation 24 (CD24) Polymorphism, Talin-1 gene expression and Hepatocellular Carcinoma prevalence in Egyptian population

Type

Research paper

Keywords

HCV, hepatocellular carcinoma, SNP, CD24, Alpha fetoprotein, Talin-1

Abstract

Introduction

Background: Hepatitis C is considered as one of the most popular diseases in Egypt. Our aim is to clarify the association between Cluster of Differentiation 24 (CD24) polymorphism, Talin-1 gene expression and the prevalence of hepatocellular carcinoma in Egyptian Hepatitis C virus patients.

Material and methods

The link between CD24 polymorphism rs8734 and the prevalence of hepatocellular carcinoma was assessed between 200 control subjects and 400 hepatitis C virus patients (HCV), patients were classified as follows; 200 patients with HCV and 200 with HCV and hepatocellular carcinoma (HCC) by histopathological assessment and PCR-restriction fragment length polymorphism (PCR-RFLP).

Results

Results: The hepatitis c patients with HCC showed significant increase in alpha fetoprotein (AFP) and Talin-1 gene expression compared to patients with HCV as well as in healthy volunteers. Furthermore, the frequencies of CD 24 170 CT/TT genotype were significantly higher in HCV patients without complications (60%) when compared to CC genotype (40%) (OR= 6 at $X^2= 14.41$, $P = 0.0007$), and in HCV with HCC patients (90%) when compared to CC genotype (10%) (OR= 36 at $X^2= 14.41$, $P = 0.0007$).

Conclusions

These data suggests that CD24 genetic polymorphism rs8734 and Talin-1 gene expression may be a significant determinant for the prevalence of hepatocellular carcinoma in HCV patients.

**Association between Cluster of Differentiation 24 (CD24)
Polymorphism, Talin-1 gene expression and Hepatocellular
Carcinoma prevalence in Egyptian population**

**Mohamed M Hafez¹, Lina Jamil M. Abdel-Hafez², Lubna Jamil³, Sherine M.
Ibrahim^{*4}**

¹Department of Biochemistry, Faculty of Pharmacy, Ahram Canadian University (ACU), October 6 City, Giza, Egypt.

²Department of Microbiology and Immunology, Faculty of Pharmacy, October 6 University (O6U), October 6 City, Giza, Egypt.

³Department of Histology, Faculty of Medicine, October 6 University (O6U), October 6 City, Giza, Egypt.

^{*4}Department of Biochemistry, Faculty of Pharmacy, Modern Sciences and Arts University (MSA), October 6 City, Giza, Egypt.

**Mohamed M Hafez: mohkhalifa2000@yahoo.com. TEL. +201061974242 ORCID:
0000-0003-0403-8033**

**Lina Jamil M. Abdel-Hafez: lina.jamil@ymail.com. TEL. +201147405051
ORCID: 0000-0001-8830-5877**

**Lubna Jamil: drlubnajamil1978@gmail.com. TEL. +201156515858 ORCID:
0000-0003-1078-7567**

***Sherine M. Ibrahim: sherinemahmoud@msa.eun.eg. TEL. +201223798354
ORCID: 0000-0002-2755-1641**

*** Corresponding Author:**

Sherine M. Ibrahim, Ph.D.

**Faculty of Pharmacy, Modern Sciences and Arts University, October 6 City,
Cairo, Egypt, postal code: 12566.**

Email: sherinemahmoud@msa.eun.eg

Telephone: +201223798354

Abstract

Background: Hepatitis C is considered one of the most popular diseases in Egypt. We aim is to clarify the association between Cluster of Differentiation 24 (CD24) polymorphism, Talin-1 gene expression, and the prevalence of hepatocellular carcinoma in Egyptian Hepatitis C virus patients.

Methods: The link between CD24 polymorphism rs8734 and the prevalence of hepatocellular carcinoma was assessed between 200 control subjects and 400 hepatitis C virus patients (HCV), patients were classified as follows; 200 patients with HCV and 200 with HCV and hepatocellular carcinoma (HCC) by histopathological assessment and PCR-restriction fragment length polymorphism (PCR-RFLP).

Results: The hepatitis c patients with HCC showed a significant increase in alpha -fetoprotein (AFP) and Talin-1 gene expression compared to patients with HCV as well as in healthy volunteers. Furthermore, the frequencies of CD 24 170 CT/TT genotype were significantly higher in HCV patients without complications (60%) when compared to CC genotype (40%) (OR= 6 at $X^2= 14.41$, $P = 0.0007$), and in HCV with HCC patients (90%) when compared to CC genotype (10%) (OR= 36 at $X^2= 14.41$, $P = 0.0007$).

Conclusion: These data suggest that CD24 genetic polymorphism rs8734 and Talin-1 gene expression may be a significant determinant for the prevalence of hepatocellular carcinoma in HCV patients.

Keywords: Hepatocellular carcinoma, SNP, HCV, CD24, Talin-1, alpha-fetoprotein.

Abbreviations:

HCV: Hepatitis C virus; ROS: Reactive oxygen species; HCC: Hepatocellular carcinoma; SNP: Single nucleotide polymorphism; CD24: Cluster of differentiation 24; GPI: Glycosyl phosphatidylinositol; AST: Aspartate aminotransferase; ALT: Alanine aminotransferase; ALP: Alkaline phosphate; GGT: Gamma Glutamyl-transferase; AFP: Alpha fetoprotein; ELISA: Enzyme Linked Immunosorbent assay; MS: Multiple sclerosis; UTRs: Untranslated regions; MDB: Mallory denk bodies; HSPs: Heat shock proteins; EMT: Epithelial-to-mesenchymal transition.

Introduction

Hepatocellular carcinoma (HCC) is recently the fifth most common type of malignancy with a high mortality rate globally [1]. In Egypt, HCC represents the fourth common cancer [2]. HCC is a complicated process that occurs due to multiple risks such as hepatitis C virus (HCV), hepatitis B virus (HBV), consumption of alcohol, and diabetes [3,4,5]. HCV causes acute and chronic hepatitis. If the hepatitis C virus is not treated, many pathways can lead to hepatocarcinogenesis [6]. The most popular pathway is that HCV core protein regulates gene expression that causes oxidative stress and leads to hepatocellular carcinoma (HCC) [7].

Furthermore, recent sequencing studies have revealed that genetic variations are associated with well-established risk factors in certain ethnic populations [8,9]. Genotype 4 is the most predominant genotype that present in Egypt, with subtype 4a that is considered the dominant subtype [10].

The cluster of differentiation 24 (CD24) is a signal transducer or heat-stable antigen (HSA) that can be defined as a human protein encoded by the CD24 gene. CD24 glycoprotein present at surfaces of neuroblasts and most B lymphocytes. It's a glycoprotein encoding gene expressed in B cells and on mature granulocytes. The glycosylphosphatidylinositol (GPI) anchors the encoded protein by cell surface links [11]

CD24 gene which is found on chromosome number six at position twenty-one is an alignment to genomic locations with similarity with that on chromosomes 1, 15, 20, and Y. Experimental determinations for corresponding translation and transcription of each genomic location are needed. CD24 polymorphism can affect the development risk of chronic HCV infection. CD24 P170T allele is associated with HCV infection at a higher level. Among the chronic HCV Egyptian patients, CD24 P170T allele shows recessive associate [12]. The rapid progression of hepatocellular carcinoma and liver cirrhosis in the CD24 P170T allele is significantly higher when compared with CD 24 P170C allele in HCV patients. Deletion of dinucleotide at position 1527 on CD24 can reduce the risks of chronic HCV infection [13]. However, only a few studies focused on the association between CD24 and the progression of HCC.

Talin-1 is a probable indicator for early diagnosis of cancer since its elevated degree of expression in blood samples from people with cancer was adequate to differentiate them from normal human specimens [14]. Talin-1 is important for cell adhesion and motility which is a very important factor in neoplasm metastasis and inflammation, also it is responsible for the activation of integrins which regulates the cell apoptosis and growth of tumors [15].

Here we conducted our study to emphasize the relation between virus C and hepatocellular carcinoma and to clarify the association between Cluster of Differentiation 24 (CD24) polymorphism, Talin-1 gene expression, and prevalence of hepatocellular carcinoma in Egyptian Hepatitis C virus patients.

Subjects and Methods:

Patient selection

A total of 600 Egyptian adults individuals were involved in this study.

Inclusion Criteria:

The inclusion criteria included age ≥ 30 years and positive HCV RNA tests for HCV patients. A total of 200 served as healthy controls and 400 patients diagnosed with HCV were selected in this study; they were selected from Theodor Bilharz Research Institute (TBRI). Patients were classified as follows: 200 patients diagnosed with chronic hepatitis C without liver carcinoma and 200 patients diagnosed with chronic hepatitis C with liver carcinoma. All cases were diagnosed according to histological assessment, and their clinical stage was determined according to the TNM staging system of the American Joint Committee on Cancer (AJCC) [16]. Liver cirrhosis was diagnosed according to abdominal sonography or liver biopsy. The evaluation of the participants for the hepatic function was based on patient history, medical consultation, serum hepatic function tests, and liver biopsies were done before any antiviral therapies were taken. The epidemiological factors included gender, body mass index (BMI), and age.

Exclusion Criteria:

Exclusion criteria were defined as: Patients infected with either hepatitis B virus (HBV) or HIV were excluded, patients having the background of any inflammatory diseases such as acute or chronic thyroid diseases, infections, and drug abuse. For alcohol consumption, patients were considered excluded when they have up to an average of more than 2 drinks per day and for persistent smoking habit, patients were considered excluded when the patient is smoking one cigarette per day in the latest three months.

The study was approved by Ahram Canadian University (ACU) Human Ethics Committee (PBC-2020-04). The study was carried out following the recommendations and regulations of the Declaration of Helsinki. Before participation, all medical histories of all subjects were collected and written informed consent was taken from all participants.

Blood sampling and laboratory assays

Blood samples were divided into two parts under complete aseptic conditions. The first part was added to tubes containing EDTA (1 mg/mL) to isolate and extract DNA by spin column-based genomic DNA by removing polymerase chain reaction (PCR) inhibitors as cations and proteins. The second part was taken into tubes, where serum is obtained by centrifugation at 4000 rpm for 15 min & serum kept frozen at -70°C for determination of Aspartate transaminase (AST) [17]., Alkaline phosphate (ALP), Alanine Transaminase (ALT), Gamma Glutamyl-transferase (GGT), total cholesterol (TC), and high-density lipoprotein (HDL) [18,19,20]., low-density lipoprotein cholesterol (LDL) using standard laboratory spectrophotometric methods and Serum Alfa-fetoprotein (AFP) levels was estimated using enzyme-linked immunoassay kits (Commercial kit purchased from DRG, USA) [21].

Genotyping of Cluster of Differentiation 24 (CD 24)

The cluster of Differentiation 24 (CD 24) gene amplification was performed using PCR-restriction fragment length polymorphism (PCR-RFLP). By measuring the concentration of each sample by fluorometer device as 1 ul of Qubit reagent was mixed with 199 ul Qubit buffer to form Qubit working solution then 199 ul from that working

solution mixed with 1 ul of DNA sample "from first step" in PCR tube after that the concentration was measured using Fluorometer device [22].

Accession Number: rs8734.

Quantitative Real-time PCR Assay of Talin-1:

Talin-1 gene expression was detected in Peripheral Blood Mononuclear Cells (PBMCs), these cells were obtained from peripheral blood by the Ficoll density sedimentation process. QIA amp viral RNA extraction kit was used for the extraction of total RNA. The quantification process was analyzed using TaqMan® Gene Expression assay (Applied Biosystems Inc, Foster City, CA, USA). Levels of Talin-1 expression were calculated using the threshold cycle method [23]. The following primers were used in the qRT-PCR [24]:

Talin-1 sense: 5'-TCTCCCAA ATGCCAAGAAC-3'
Anti-sense: 5'-TGGCTATTGG GGTCAGAGAC-3'
Glyceraldehyde -3- phosphate dehydrogenase (GAPDH) sense: 5'-
CCACTCC TCCACCTTTGAC-3'
Anti-sense: 5'-ACCCTGT TGCTGTAGCCA-3'

Hepatic histopathological assessment by: Hematoxylin and eosin & Masson's trichrome staining

Hematoxylin and eosin staining

Liver specimens were taken by needle biopsy from all patients with hepatitis C upon signing a written consent. Liver tissues were all treated with 10 percent neutral buffered formalin for 24 h and histo-processed. The blocks were then sliced into 3 μ thicknesses, using a rotary microtome. Parts have been stained utilizing hematoxylin & eosin stain [25].

Masson's trichrome staining

Three colors can be used for muscle staining, collagen fibers, fibrin, and erythrocytes. The fundamental principle in trichrome staining is that less porous tissue

is colored by the small dye molecule; whereas a higher molecular dye can infiltrate [26]. Hematoxylin and eosin & Masson's trichrome staining have been used for histopathological analysis. The level of steatosis, lobular inflammatory processes, and non-alcoholic steatohepatitis (NASH) was calculated using the SAF scoring system. Steatosis, Activity & Fibrosis Score System (SAF) is a non-alcoholic fatty liver disease (NAFLD) score based on histological severity. All biopsies were classified according to the SAF system, and the severity of the disease was classified as mild, moderate or severe. The SAF activity score was measured by hepatocellular ballooning and lobular inflammation. Histopathological severe disease was described as SAF activity score > 3 for bridging fibrosis or cirrhosis. The regression formula for the fibrosis severity estimation includes 6 variables: age, years, BMI: Kg/m², fasting glucose (FG)/ diabetes, Platelet count & albumin (g/dl) and AST / ALT ratio [27].

Statistical Analysis:

The demographic characteristics differences were compared using Fisher's exact test and the Mann–Whitney U test between healthy controls and HCC patients were. These data were expressed as Mean ± SEM and compared between groups by the Student's t-test. The data were analyzed by using SPSS 25 software (IBM SPSS, USA). *P values* < 0.05 were regarded as statistically significant.

Results:

Characteristics of Subjects

Since various risk factors have been related to the pathogenesis and prevalence of liver cancer such as alcohol consumption, gender, and age, we first compared the mean and SEM for the clinical data of 400 HCV patients with that from 200 normal controls (**Table 1**). There was no significant difference between the study groups in the sex distribution, BMI, or age. For the liver function tests, the levels of ALT are elevated significantly in hepatitis C patients (Mean ± SEM = 125± 20.53 IU/L) and Hepatitis C with hepatocellular carcinoma patients (Mean ± SEM = 135 ± 11.69 IU/L) when compared to healthy control subjects (Mean ± SEM = 27.8 ± 0.61 IU/L). Meanwhile, there is no significance in it between the HCV group and HCV with the HCC group at *P* < 0.05. While for GGT and ALP levels, both were significantly higher in all HCV

patients than controls. Also, there is a significant increase in it between HCV with HCC group and HCV group at $P < 0.05$. Additionally, the serum AFP level in Hepatitis C with hepatocellular carcinoma patients showed the highest significant increase to 4570 ± 294 ng/mL compared to the healthy control group 6.6 ± 0.8 ng/mL ($P \leq 0.05$). Additionally, we observed that tobacco smoking and alcohol consumption have a high tendency to affect the prevalence of HCC (**Table 1**). The power of study was calculated as; Total sample size:600, Number of groups:3, Effect size :0.15, critical F: 3, and Power:0.9

Histopathological Examination

Figure (1) showed a photomicrograph in the control human liver section showing normal hepatic lobules architecture. While in **figure (2)** the histopathological sections revealed human liver sections showing (A and B): Mallory-Denk bodies & microvesicular steatosis (C): a human liver section with NASH was demonstrated showing Peri-sinusoidal fibrosis (Masson's trichrome) with no CD4 expression. On the contrary, histology analysis in **figure (3)** indicates that the tumors in hepatitis c patients with complications group are malignant HCC, with the expression of CD4 on the inflammatory cells which are determined by its heterogeneous and large nuclei, cancer cells are also characterized by double nuclei, showing: (A): Masses of malignant cells with frequent mitosis, hyperchromatic nuclei & trabecular growth pattern (HX&E 100x). (B): Pseudo glandular growth pattern (HX&E 100X) (C): Loss of architecture, cellular degeneration & Solid growth pattern (HX&E 200x). (D): Loss of architecture with dilated central vein & giant cell formation. These data focused on the correlation between CD24 gene variation and the rapid development of hepatocellular carcinoma.

Genotype distribution and allele frequencies of CD 24 170 C/T polymorphism in hepatitis C patients' groups versus the control group

The frequencies of CD 24 170 CT/TT genotype were significantly higher in hepatitis C patients without complications (60%) when compared to CC genotype (40%) with odds ratio= 6 at $X^2 = 14.41$, $P = 0.0007$, as well as, the frequencies of CD 24 170 CT/TT genotype were significantly higher in hepatitis C patients with HCC (90%) when compared to CC genotype (10%) with odds ratio= 36 at $X^2 = 14.41$, $P = 0.0007$ (**Table 2**). No significant differences were found between the levels of AFP,

AST, ALT, or GGT and the CD 24 170 C/T polymorphisms in HCC patients (**Table 3**). CD24 polymorphism correlated with prognosis in HCC patients which is estimated by its correlation with tumor status T1+T2 at $P < 0.0001$ in figure (4). Additionally, Kaplan-Meier curve was conducted to estimate the overall survival rate between the different genotypes of CD24 polymorphism of the recipients at $P < 0.0001$ as estimated in figure (5).

Talin-1 Gene Expression

Table (4) showed a 1.6-fold significant increase in of Talin-1 expression in HCV patients (Mean \pm SEM = 8.07 ± 0.12 & 14.27 ± 0.12 respectively) when compared to control group (Mean \pm SEM = 6.15 ± 0.1) at $P < 0.0001$. Moreover, there was a 1.3 significant increase of Talin-1 expression in HCV with HCC patients (Mean \pm SEM = 10.27 ± 0.12) compared to HCV patients (Mean \pm SEM = 8.07 ± 0.12) at $P < 0.0001$ as shown in figure (6). A significant correlation between CD24 polymorphism genotypes and talin-1 gene expression is clearly represented among our groups at $P < 0.0001$ in figure (7). **To evaluate the diagnostic value of Talin-1 gene expression, we used ROC methods to calculate the sensitivity and specificity as shown in figure 8. The AUC of Talin-1 gene expression between the patients with HCV and the HCC with those with HCV was 0.9 (95% confidence interval, 0.52-0.69; P = 0.009) for predicting the risk of developing cancer, indicating that Talin-1 gene expression had significant accuracy as a predictor for cancer prevalence risk.**

SAF Scores to Biopsies of HCV Patients

Table (5 & 6) showed the most frequent diagnosis of hepatitis C patients. The baseline diagnosis with NASH was concordant with the reference classification which states that less than -1.455 indicates the absence of fibrosis (F0-F2), between -1.455 & 0.675 indicates indeterminate score while more than 0.675 indicates the presence of fibrosis (F3-F4). Table (5) showed the most frequent diagnosis among the two pathologists in HCV patients with hepatocellular carcinoma. 40 out of 70 (57%) diagnosed with NASH were concordant with the reference classification. While in Table (6); 34 out of 70 (49%) of HCV patients without hepatocellular carcinoma were diagnosed with NASH with a full agreement between reference and baseline classification.

Table (7) showed the distribution frequency of CD24 gene variation among the 400 patients with HCV with their clinical status and histological examinations. The patients with HCV were evaluated to understand the effect of CD 24 170 CT/TT genotypes on the clinical stage, lymph node involvement, distant metastasis, vascular invasion, and liver cirrhosis. There was a significant difference in the effect of CD 24 170 CT/TT genotype on lymph node involvement, distant metastasis and liver cirrhosis as well.

Discussion

In our study, we investigated the association between CD24 polymorphisms and the prevalence of HCC in HCV patients. Hepatocellular carcinoma is the third most common cause of death due to cancer worldwide [28]. In Egypt, hepatocellular carcinoma represents around 1.68% of the total malignancies and 11.75% of all digestive organs' malignancies and metabolic syndrome diseases [29]. The distribution analysis of the CD24 genotypes involved 400 HCV patients and 200 normal controls indicating the frequencies of CD 24 170 CT/TT genotype were significantly higher in HCV patients without complications (60%) when compared to CC genotype (40%) with odds ratio= 6 at $X^2= 14.41$, $P = 0.0007$, as well as, the frequencies of CD 24 170 CT/TT genotype were significantly higher in HCV with HCC (90%) when compared to CC genotype (10%) with odds ratio= 36 at $X^2= 14.41$, $P = 0.0007$.

Moreover, among the chronic HCV Egyptian patients, CD24 P170T allele shows a strong association with the rapid development of hepatocellular carcinoma and liver cirrhosis when compared with CD 24 P170 C allele. In contrast, deletion of dinucleotide at position 1527 on CD24 can reduce the development of chronic HCV infection [30]. In addition, CD24 polymorphisms may increase the genetic susceptibility factor for HBV infection. This was investigated by a recent study which included 609 HBV patients and 383 healthy controls, this study showed an increased risk of prevalence of HBV infection in patients with the P170T/T genotype when compared with P170C/T and P170C/C genotypes using the logistic regression model [31].

An important explanation for how the CD24 SNP affects the risk of development of chronic HCV infection. Our previous results have suggested that the P170T allele which is expressed at a higher level than P170C encodes a certain protein, which is responsible for the progression of chronic HCV infection by affecting the efficiency of cleavage of posttranslational GPI. These results agree with the study of [32] which supports the idea that P170T allele affects the progression of chronic HCV infection through posttranslational mechanisms. Given the fact that CD24 is expressed mainly in the neuronal cells and hematopoietic, Huang and Hsu³¹ showed that there are many other tumor cells that have been shown to increase the expression of CD24 mainly in liver tumors [21]. Another study by [33] also suggested that CD24 SNPs are a prognostic marker for hepatic carcinoma. In this regard, another study confirmed that during liver carcinogenesis CD24 is highly expressed in the liver progenitor cells [34].

Furthermore, CD24 polymorphism may affect the immune/inflammatory response through T-cell activation. T-cell-mediated inflammation is one of the important mechanisms for the prevalence of HCC in HBV infected mice, it also affects the production of cytokines from liver necrotic cells [35] CD24 P170 T/T is a higher cell-surface genotype than P170 C/T or P170 C/C genotypes, which increases the rapid progression and risk of multiple sclerosis (MS). While dinucleotide deletion in 3 untranslated regions (UTRs) is associated with protection from systemic lupus erythematosus and MS, as that deletion reduces the stability of CD24 messenger RNA [36].

Talin-1 serves an important character in the stimulation of integrin. Especially, the sensitivity of Talin-1 for cancer diagnosis was stronger than that of AFP in Egyptian Patients with HCC [37]. Obviously, these results indicate that Talin-1 is a possible diagnostic indicator for HCC. This was also investigated by another recent study which was performed on 90 Egyptian subjects showing that talin-1 is involved in the carcinogenesis of HCC [14] Even so, whether Talin-1 stimulated HCC proliferation and metastases were still unknown, and the Talin-1 role in HCC proliferation remained under investigation [38].

Talin-1 has been shown to facilitate HCC progression by inhibiting the activity of apoptosis considerations and the anti-apoptotic BCL₂ members [39] It also encourages HCC metastasis by inducing the release of mesenchymal Epithelial-to-

mesenchymal transition (EMT) markers and by reducing the activity of epithelial molecules. Talin-1 can stimulate HCC expansion and metastases through the regulation of electrical cell signaling and function as a promising bioelectricity target therapeutically [40]. In our research, the histological characteristics of hepatocellular carcinoma are prominent acinar patterns, mitotic activity, pseudo glandular or acinar & compact or solid patterns. Septae are observed & giant cells are also seen. The occurrence of liver biopsy steatosis in HCV patients is more expressed compared to other liver diseases such as autoimmune hepatitis and chronic hepatitis B. Steatosis is suggested to be 2.5 times more common in HCV patients relative to the normal community. Macrovesicular steatosis occurring in HCV patients is often distributed in the periportal and non-centrilobular regions, which are most frequently seen in NAFLD. All this implies that HCV can directly induce steatosis in these patients [41].

In summary, our study suggested that CD24 polymorphism P170 CT/TT may affect both the prevalence of chronic HCV infection and HCC. Moreover, Talin-1 gene expression was shown to facilitate HCC progression by reducing the activity of epithelial cells and through regulation of electrical cell signaling and inhibition of apoptosis.

Conclusion:

The expression of CD24 polymorphism and talin-1 gene outside the hematopoietic cells recently raised an interesting concern as a promising prognostic marker for progression of chronic HCV infection and HCC with better accuracy than serum AFP due to their high correlation with invasion and malignant growth of hepatocytes and the immune/inflammatory response in the liver tissue.

Glossary:

- **Single Nucleotide Polymorphism (SNP):** is a substitution of a single nucleotide that happens at a particular position in the genome.
- **CD24:** is a sialoglycoprotein which expressed at the B lymphocytes' surface and differentiating neuroblasts.
- **HCV:** is a type of viral liver disease that happens as acute or chronic.

- **Hepatocellular Carcinoma:** is a liver malignancy which found predominantly in patients with significant chronic liver disorder and cirrhosis.

Declarations

Authors' contributions: Lina Jamil and M. Abdel-Hafez participated in the study design, practical work especially the molecular biology as well as collection of data and manuscript preparation. Sherine M. Ibrahim and Mohamed M. Hafez participated in practical work, data analysis and preparation of manuscript. Lubna Jamil participated in the histopathological work. All authors revised and approved the final manuscript.

Acknowledgements: We thank Theodor Bilharz Research Institute (TBRI) for helping in samples' collection.

Formatting of funding sources: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical approval: The study protocol was approved by the Ethics Committee of the Faculty of Pharmacy, Ahrum Canadian University (ACU) **Human Ethics Committee (PBC-2020-04).**

Conflict of interest: The authors declare that they have no conflict of interest.

Consent for publication

The signed Consent ensures that the Publisher has the Author's permission to publish the relevant Contribution.

Data Availability Statement: All data generated or analysed during this study are included in this published article.

Accession Number: rs8734.

References

1. Shawky E, M. El-Beih N, El-Husseyeny E, El-Ahwany E, Hassan M, Zoheiry M. Effects of free and nanoparticulate curcumin on chemically induced liver carcinoma in an animal model. *Arch Med Sci* 2021;17(1):218–227
2. Abd-Elsalam S, Elwan N, Soliman H, et al. Epidemiology of liver cancer in Nile delta over a decade: a single-center study. *South Asian J Cancer* 2018; 7:24.
3. Sherman, M. Hepatocellular carcinoma: Epidemiology, surveillance, and diagnosis. *Semin. Liver Dis* 2010; 30: 3–16.
4. Forner A, Lovet JM, Bruix J. Hepatocellular carcinoma. *Lancet* 2012; 379: 1245–1255.
5. Gao J, Xie L, Yang WS, et al. Risk factors of hepatocellular carcinoma—Current status and perspectives. *Asian Pac. J. Cancer Prev* 2012; 13: 743–752.
6. Sharma SD. Hepatitis C virus: molecular biology & current therapeutic options. *Indian J Med Res* 2010; 131: 17- 34.
7. Gottwein JM, Scheel TK, Jensen TB, et al. Development and characterization of hepatitis C virus genotype 1-7 cell culture systems: role of CD81 and scavenger receptor class B type I and effect of antiviral drugs. *Hepatology* 2009; 49 (2): 364- 377.
8. Miki D, Ochi H, Hayes CN, Aikata H, Chayama K. Hepatocellular carcinoma: Towards personalized medicine. *Cancer Sci* 2012; 103: 846–850.
9. Nahon, P, Zucman-Rossi, J. Single nucleotide polymorphisms and risk of hepatocellular carcinoma in cirrhosis. *J. Hepatol* 2012; 57: 663–674.
10. Jin, F, Xiong, W, Jing JC, Feng Z, Qu LS, Shen X. Evaluation of the association studies of single nucleotide polymorphisms and hepatocellular carcinoma: A systematic review. *J. Cancer Res* 2011; 137: 1095–1104.
11. Simmonds P, Bukh J, Combet C, Deléage G, et al. Consensus proposals for a unified system of nomenclature of hepatitis C virus genotypes. *Hepatology* 2005; 42 (4): 962- 973.

12. Siegel R, Desantis C, Jemal A. Colorectal cancer statistics. *Cancer J Clin* 2014; 64 (2): 104- 117.
13. Braliou GG, Pantavou KG, Kontou PI, Bagos PG. Polymorphisms of the CD24 gene are associated with risk of multiple sclerosis: A Meta-Analysis. *Int J Mol Sci* 2015; 16 (6): 12368- 12381.
14. Austen K, Ringer P, Mehlich A, et al. Extracellular rigidity sensing by talin isoform-specific mechanical linkages. *Nat Cell Biol.* 2015; 17(12):1597-1606.
15. Mashaly A, Aya H, et al. Diagnostic and prognostic value of Talin-1 and Midkine as tumor markers in hepatocellular carcinoma in Egyptian patients. *Asian Pacific journal of cancer prevention* 2018; 1503.
16. Vauthey JN, Lauwers G, Esnaola N, et al. Simplified staging for hepatocellular carcinoma. *J. Clin. Oncol* 2002; 20: 1527–1536
17. Kim HJ, Kim SY, Shin SP, et al. Immunological measurement of aspartate/alanine aminotransferase in predicting liver fibrosis and inflammation. *Korean J Intern Med* 2019.
18. Karmen A. A note on the spectrometric assay of glutamic-oxalacetic transaminase in human blood serum. *J Clin Invest* 1955; 34 (1): 131- 133.
19. Yoon S, Lim N, Ha S, et al. The human cervical cancer oncogene protein is a biomarker for human hepatocellular carcinoma. *Cancer Res* 2004; 64: 5434-5441.
20. Hall P, Cash J. “What is the Real Function of the Liver Function Tests?”. *Ulster Med J* 2012; 81 (1): 30-36.
21. Huang G, Yang L, Lu W. Expression of hypoxia-inducible factor and vascular endothelial growth factor in hepatocellular carcinoma: Impact on neovascularization and survival. *World J Gastroenterol* 2005; 11: 1705-1708.
22. Kawada N. Evolution of hepatic fibrosis research. *Hepatol Res* 2011; 41 (3): 199- 208.
23. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2 Delta C(T) Method. *Methods* 2001; 25(4): 402-408.
24. Sun J, Fang K, Shen H, Qian Y. MicroRNA-9 is a ponderable index for the prognosis of human hepatocellular carcinoma. *Int J Clin Exp Med* 2015; 8(10): 17748-17756.

25. Alwahaibi NY, Alkhatri AS, Kumar JS. Hematoxylin and eosin stain shows a high sensitivity but sub-optimal specificity in demonstrating iron pigment in liver biopsies. *Int J Appl Basic Med Res* 2015;5(3):169-171.
26. Forlano R, Mullish BH, Giannakeas N, et al. High Machine Learning-Based Quantification of Steatosis, Inflammation, Ballooning, and Fibrosis in biopsies from Patients with Nonalcoholic Fatty Liver Disease. *Clin Gastroenterol Hepatol.* 2020;18(9):2081-2090.
27. Munteanu M, Tiniakos D, Anstee Q, et al. Diagnostic performance of Fibro-Test, Steato-Test and Acti -Test in patients with NAFLD using the SAF score as histological reference. *Aliment Pharmacol Ther* 2016; 44:877–889.
28. Liu J, Chen L, Pan J, Chen M, Zhou J, Zhou F, Chen P, Song Y. Comprehensive analysis of key lncRNAs in HCV-positive hepatocellular carcinoma. *Arch Med Sci* 2021;17(1):142–151
29. Sherine M. Ibrahim and Afaf A. Bastawy. The Relevance of Single-nucleotide Polymorphism +62 G>A to the Expression of Resistin Gene Affecting Serum Resistin Levels in Metabolic Syndrome in the Egyptian Population. *Curr Pharm Biotechnol.* 2020;21(7):626-634.
30. Wang L, Lin S, Rammohan KW, et al. A dinucleotide deletion in CD24 confers protection against autoimmune diseases. *PLoS Genet* 2007; 3 (4): e49.
31. Li D, et al. CD24 polymorphisms affect risk and progression of chronic hepatitis B virus infection. *Hepatology* 2009,19: 735-742.
32. Robert F, Pelletier J. Exploring the Impact of Single-Nucleotide Polymorphisms on Translation. *Front Genet* 2018; 9: 507.
33. Kristiansen G, Machado E, Bretz N, et al. Molecular and clinical dissection of CD24 antibody specificity by a comprehensive comparative analysis. *Lab Invest* 2010; 90 (7): 1102- 1116.
34. Ochsner SA, Strick-Marchand H, Qiu Q, et al. Transcriptional profiling of bipotential embryonic liver cells to identify liver progenitor cell surface markers. *Stem Cells* 2007; 25:2476-2487
35. Mihm S. Hepatitis C virus, diabetes and steatosis: clinical evidence in favor of a linkage and role of genotypes. *Dig Dis* 2010; 28 (1): 280- 284.
36. Mirza S, Siddiqui AR, Hamid S, Umar M, Bashir S. Extent of liver inflammation in predicting response to interferon α & Ribavirin in chronic hepatitis C patients: a cohort study. *BMC Gastroenterol* 2012; 14: 12-71.

37. Youns MM, Abdel Wahab AH, Hassan ZA, Attia MS. Serum talin-1 is a potential novel biomarker for diagnosis of hepatocellular carcinoma in Egyptian patients. *Asian Pac J Cancer Prev.* 2013; 14(6):3819-23.
38. Bostanci O, Kemik O, Kemik A, etal. A novel screening test for colon cancer: Talin-1. *Eur Rev Med Pharmacol Sci* 2014; 18(17):2533-2537.
39. Sakamoto S, McCann RO, Dhir R, Kyprianou N. Talin1 promotes tumor invasion and metastasis via focal adhesion signaling and anoikis resistance. *Cancer Res.* 2010; 70: 1885-1895.
40. Chen P, Zheng X, Zhou Y, Xu Y, Zhu L, Qian Y. Talin-1 interaction network promotes hepatocellular carcinoma progression. *Oncotarget* 2017;21;8(8):13003-13014.
41. Zampino R, Marrone A, Rinaldi L, etal. Endotoxemia contributes to steatosis, insulin resistance and atherosclerosis in chronic hepatitis C: the role of pro-inflammatory cytokines and oxidative stress. *Infection.* 2018;46(6):793-799.

Preprint

Table 1: Clinical and Hemodynamic Characteristics of Participants.

Variables	Control	HCV	HCV with HCC	P value
Sex	200 (35 M / 35 F)	200 (30 M /40 F)	200 (35M / 35F)	0.01
Age (years)	40.51 ± 0.66	41.38 ± 0.55	40.38 ± 0.55	0.05
BMI (Kg/m²)	23 ± 0.22	24 ± 0.26	24 ± 0.12	0.01
Waist	0.75	0.82	0.83	0.01
Waist/hip ratio	0.83	0.85	0.84	0.01
Cigarette smoking				0.01
No	170	55	45	
Yes	30	145	155	
Alcohol consumption				0.001
No	55	26	20	
Yes	15	44	50	
Tumor status				0.001
T1+T2			128(64%)	
T3+T4			72(36%)	
SBP (mm Hg)	120.3 ± 8.44	140 ± 7.32 ^{a,b}	145 ± 7.32 ^{a,b}	0.01
DBP (mm Hg)	70.5 ± 5.41	85± 4.32 ^{a,b}	95± 4.32 ^{a,b}	0.01
Serum TAG (mg/dL)	111.6 ± 2.2	171.9 ± 4.05 ^a	175.9 ± 4.05 ^a	0.001
Serum TC (mg/dL)	150.3 ± 1.70	230.7 ± 3.09 ^a	228.7 ± 3.09 ^a	0.001
Serum HDL-C (mg/dL)	56.53 ± 0.57	29.71 ± 0.92 ^a	30.71 ± 0.92 ^a	0.01
Serum LDL-C (mg/dL)	95.37 ± 1.81	150.6 ± 3.4 ^a	155.6 ± 3.4 ^a	0.01
GGT (IU/L)	21.5 ± 0.77	85.3 ± 0.66 ^a	96.1 ± 0.69 ^{ab}	0.01
ALP (IU/L)	60 ± 2.78	192 ± 1.56 ^a	176.3±0.85 ^{ab}	0.001
AST (IU/L)	20.9 ± 0.45	120.5 ± 14.57 ^a	131 ± 10.77 ^{ab}	0.001
ALT (IU/L)	27.8 ± 0.61	125± 20.53 ^a	135 ± 11.69 ^a	0.001
Serum AFP (ng/mL)	6.6 ± 0.8	2800.9 ± 110 ^a	4570 ± 294 ^{ab}	0.01

Abbreviations: HCV, hepatitis C virus; BMI, body mass index; TAG, triacylglycerol; TC, total cholesterol; HDL-C, high-density lipoprotein; LDL-C, low-density lipoprotein cholesterol. ALP; Alkaline phosphatase test. AST; Aspartate aminotransferase. ALT; Alanine aminotransferase. GGT; gamma glutamyl- transferase. AFP, alpha-fetoprotein.

Data are given as mean + SEM

^a Significant difference from control group at $P < 0.05$

^b Significant difference from HCV group at $P < 0.05$

Preprint

Table 2: Difference in genotype frequency of CD 24 SNP 170 between all studied groups

Groups	N	Genotype Frequency		OR	95 % CI
		CC	CT + TT		
Control	200	160 (80 %)	40 (20 %)		
HCV	200	80 (40 %)	120 (60 %)	6	0.88- 13.83
HCV with HCC	200	20 (10 %)	180 (90 %)	36	2.07- 30.89
$X^2 = 14.41, P=0.0007^*$					

X²: Chi square, OR: Odd ratio, CI: Confidence interval.

*Statistically significant difference at p< 0.05 using Chi square test.

Preprint

Table 3: Association of CD 24 C/T SNP 170 genotypic frequencies with HCC laboratory status.

Characteristic	α-Fetoprotein ^a (ng/mL)	AST ^a (IU/L)	ALT ^a (IU/L)	GGT ^a
CD 24 C/T SNP 170				
CC	2800.2 ± 110	125.1 ± 20.6	120.2 ± 14	85 ± 0.66
CT + TT	4570 ± 294	135.2 ± 11.6	131.3 ± 10	96 ± 0.69
p value	0.448	0.537	0.501	0.545

Mann-Whitney U test was used between two groups.

a Mean ± S.E.

*** p value < 0.05 as statistically significant.**

Table 4: Gene Expression of Talin-1 in hepatitis C virus patients (HCV patients) and hepatitis C virus with hepatocellular carcinoma patients (HCV with HCC patients) when compared to control group at P < 0.05:

Group	Talin-1 gene expression
Control	6.15 ± 0.1
HCV	8.07 ± 0.12 ^a
HCV with HCC	10.27 ± 0.12 ^{ab}

a Significant when compared to control group at P < 0.05

b Significant when compared to control group at P < 0.05

Preprint

Table 5: Evaluation of liver Biopsies by the Liver Pathologists in HCV patients with hepatocellular carcinoma

Case	Reference classification	Algorithmic classification	SAF	Disease severity
1	NASH	NASH	S2A4F1	Significant
2	steatosis	steatosis	S1A1F1	Mild
3	NASH	NASH	S1A4F3	Significant
4	NASH	NASH	S3A1F2	Significant
5	steatosis	steatosis	S3A1F0	Mild
6	NASH	NASH	S3A3F2	Significant
7	NASH	NASH	S2A4F1	Significant
8	steatosis	steatosis	S1A1F1	Mild
9	NASH	NASH	S3A1F2	Significant
10	NASH	NASH	S2A4F1	Significant
11	NASH	NASH	S3A1F2	Significant
12	NASH	NASH	S2A4F1	Significant
13	steatosis	steatosis	S1A1F1	Mild
14	NASH	NASH	S3A3F2	Significant
15	NASH	NASH	S2A4F1	Significant
16	steatosis	steatosis	S1A1F1	Mild
17	steatosis	steatosis	S1A1F1	Mild
18	NASH	NASH	S3A4F3	Significant
19	NASH	NASH	S3A3F2	Significant
20	NASH	NASH	S2A4F1	Significant
21	NASH	NASH	S2A4F1	Significant
22	steatosis	steatosis	S1A1F1	Mild
23	steatosis	steatosis	S1A0F0	Mild
24	steatosis	steatosis	S1A1F1	Mild
25	NASH	NASH	S3A3F2	Significant
26	NASH	NASH	S3A1F2	Significant
27	NASH	NASH	S3A1F2	Significant
28	NASH	NASH	S2A4F1	Significant
29	steatosis	steatosis	S1A0F0	Mild
30	steatosis	steatosis	S1A1F1	Mild
31	steatosis	steatosis	S1A0F0	Mild
32	NASH	NASH	S3A3F2	Significant
33	NASH	NASH	S3A1F2	Significant
34	NASH	NASH	S3A1F2	Significant
35	NASH	NASH	S2A4F1	Significant
36	NASH	NASH	S3A1F2	Significant
37	steatosis	steatosis	S1A1F1	Mild
38	NASH	NASH	S3A1F2	Significant
39	NASH	NASH	S3A1F2	Significant
40	NASH	NASH	S2A4F1	Significant
41	steatosis	steatosis	S3A1F0	Mild
42	steatosis	steatosis	S1A1F1	Mild
43	steatosis	steatosis	S1A0F0	Mild
44	NASH	NASH	S3A1F2	Significant
45	steatosis	steatosis	S3A1F0	Mild
46	steatosis	steatosis	S1A1F1	Mild
47	NASH	NASH	S3A3F2	Significant
48	NASH	NASH	S3A3F2	Significant
49	NASH	NASH	S2A4F1	Significant
50	steatosis	steatosis	S3A1F0	Mild
51	NASH	NASH	S3A3F2	Significant
52	steatosis	steatosis	S1A0F0	Mild
53	steatosis	steatosis	S1A1F1	Mild
54	NASH	NASH	S3A3F2	Significant
55	NASH	NASH	S2A4F1	Significant
56	NASH	NASH	S3A3F2	Significant
57	steatosis	steatosis	S2A1F1	Mild
58	steatosis	steatosis	S1A1F1	Mild
59	NASH	NASH	S3A1F2	Significant
60	steatosis	steatosis	S2A1F1	Mild
61	steatosis	steatosis	S2A1F1	Mild
62	NASH	NASH	S2A4F1	Significant
63	NASH	NASH	S3A3F2	Significant
64	NASH	NASH	S3A1F2	Significant
65	NASH	NASH	S2A4F1	Significant
66	steatosis	steatosis	S1A1F1	Mild
67	steatosis	steatosis	S2A1F1	Mild
68	steatosis	steatosis	S1A1F1	Mild
69	NASH	NASH	S3A1F2	Significant
70	NASH	NASH	S2A4F1	Significant
71	NASH	NASH	S3A1F2	Significant
72	NASH	NASH	S2A4F1	Significant

*Reference interpretation: is the initial evaluation done by pathologist using accepted criteria, algorithm and SAF score.

*Algorithmic Classification: (Steatosis vs. NASH) after using the algorithm.

* Based on SAF score: mild for A < 2 and F < 2 or significant for A > 2 and/or F > 2.

73	steatosis	steatosis	S1A0F0	Mild
74	steatosis	steatosis	S1A1F1	Mild
75	steatosis	steatosis	S1A1F1	Mild
76	steatosis	steatosis	S1A1F1	Mild
77	NASH	NASH	S3A1F2	Significant
78	steatosis	steatosis	S1A1F1	Mild
79	steatosis	steatosis	S1A1F1	Mild
80	steatosis	steatosis	S1A1F1	Mild
81	steatosis	steatosis	S1A0F0	Mild
82	NASH	NASH	S3A1F2	Significant
83	steatosis	steatosis	S3A1F0	Mild
84	steatosis	steatosis	S1A1F1	Mild
85	NASH	NASH	S3A3F2	Significant
86	NASH	NASH	S3A3F2	Significant
87	NASH	NASH	S2A4F1	Significant
88	steatosis	steatosis	S3A1F0	Mild
89	NASH	NASH	S3A3F2	Significant
90	steatosis	steatosis	S1A0F0	Mild
91	steatosis	steatosis	S1A1F1	Mild
92	NASH	NASH	S3A3F2	
93	NASH	NASH	S2A4F1	Significant
94	NASH	NASH	S3A3F2	Significant
95	steatosis	steatosis	S2A1F1	Mild
96	steatosis	steatosis	S1A1F1	Mild
97	NASH	NASH	S3A1F2	Significant
98	steatosis	steatosis	S2A1F1	Mild
99	steatosis	steatosis	S2A1F1	Mild
100	NASH	NASH	S2A4F1	Significant
101	NASH	NASH	S3A3F2	Significant
102	NASH	NASH	S3A1F2	Significant
103	NASH	NASH	S2A4F1	Significant
104	NASH	NASH	S3A1F2	Significant
105	steatosis	steatosis	S1A1F1	Mild
106	steatosis	steatosis	S1A1F1	Mild
107	NASH	NASH	S3A4F3	Significant
108	NASH	NASH	S3A3F2	Significant
109	NASH	NASH	S2A4F1	Significant
110	NASH	NASH	S3A1F2	Significant
111	NASH	NASH	S2A4F1	Significant
112	steatosis	steatosis	S1A0F0	Mild
113	NASH	NASH	S2A4F1	Significant
114	steatosis	steatosis	S1A0F0	Mild
115	NASH	NASH	S3A1F2	Significant
116	NASH	NASH	S3A1F2	Significant
117	steatosis	steatosis	S2A1F1	Mild
118	steatosis	steatosis	S2A1F1	Mild
119	NASH	NASH	S2A4F1	Significant
120	NASH	NASH	S3A3F2	Significant
121	NASH	NASH	S3A1F2	Significant
122	NASH	NASH	S2A4F1	Significant
123	NASH	NASH	S3A1F2	Significant
124	steatosis	steatosis	S1A1F1	Mild
125	steatosis	steatosis	S1A1F1	Mild
126	NASH	NASH	S3A4F3	Significant
127	NASH	NASH	S3A3F2	Significant
128	NASH	NASH	S2A4F1	Significant
129	NASH	NASH	S3A1F2	Significant
130	NASH	NASH	S2A4F1	Significant
131	steatosis	steatosis	S1A0F0	Mild
132	NASH	NASH	S2A4F1	Significant
133	steatosis	steatosis	S1A0F0	Mild
134	steatosis	steatosis	S1A0F0	Mild
135	steatosis	steatosis	S1A0F0	Mild
136	NASH	NASH	S3A1F2	Significant
137	NASH	NASH	S2A4F1	Significant
138	steatosis	steatosis	S1A0F0	Mild
139	steatosis	steatosis	S1A0F0	Mild
140	steatosis	steatosis	S1A0F0	Mild
141	NASH	NASH	S3A1F2	Significant
142	NASH	NASH	S2A4F1	Significant

143	steatosis	steatosis	S1A0F0	Mild
144	steatosis	steatosis	S1A0F0	Mild
145	steatosis	steatosis	S1A1F1	Mild
146	steatosis	steatosis	S1A0F0	Mild
147	NASH	NASH	S3A3F2	Significant
148	NASH	NASH	S2A4F1	Significant
149	NASH	NASH	S3A1F2	Significant
150	NASH	NASH	S2A4F1	Significant
151	steatosis	steatosis	S1A0F0	Mild
152	NASH	NASH	S2A4F1	Significant
153	steatosis	steatosis	S1A0F0	Mild
154	steatosis	steatosis	S1A0F0	Mild
155	steatosis	steatosis	S1A1F1	Mild
156	steatosis	steatosis	S1A0F0	Mild
157	NASH	NASH	S3A3F2	Significant
158	NASH	NASH	S2A4F1	Significant
159	NASH	NASH	S3A1F2	Significant
160	NASH	NASH	S2A4F1	Significant
161	steatosis	steatosis	S1A0F0	Mild
162	NASH	NASH	S2A4F1	Significant
163	steatosis	steatosis	S1A0F0	Mild
164	steatosis	steatosis	S1A0F0	Mild
165	steatosis	steatosis	S1A1F1	Mild
166	steatosis	steatosis	S1A0F0	Mild
167	steatosis	steatosis	S1A0F0	Mild
168	steatosis	steatosis	S1A1F1	Mild
169	steatosis	steatosis	S1A0F0	Mild
170	NASH	NASH	S3A3F2	Significant
171	NASH	NASH	S2A4F1	Significant
172	NASH	NASH	S3A1F2	Significant
173	NASH	NASH	S2A4F1	Significant
174	steatosis	steatosis	S1A0F0	Mild
175	NASH	NASH	S2A4F1	Significant
176	steatosis	steatosis	S1A0F0	Mild
177	NASH	NASH	S3A1F2	Significant
178	steatosis	steatosis	S2A1F1	Mild
179	steatosis	steatosis	S2A1F1	Mild
180	NASH	NASH	S2A4F1	Significant
181	NASH	NASH	S3A3F2	Significant
182	NASH	NASH	S3A1F2	Significant
183	NASH	NASH	S2A4F1	Significant
184	NASH	NASH	S3A1F2	Significant
185	steatosis	steatosis	S1A1F1	Mild
186	steatosis	steatosis	S1A1F1	Mild
187	NASH	NASH	S3A4F3	Significant
188	NASH	NASH	S3A3F2	Significant
189	NASH	NASH	S2A4F1	Significant
190	NASH	NASH	S3A1F2	Significant
191	NASH	NASH	S2A4F1	Significant
192	steatosis	steatosis	S1A0F0	Mild
193	NASH	NASH	S2A4F1	Significant
194	steatosis	steatosis	S1A0F0	Mild
195	NASH	NASH	S3A1F2	Significant
196	NASH	NASH	S3A1F2	Significant
197	steatosis	steatosis	S2A1F1	Mild
198	steatosis	steatosis	S2A1F1	Mild
199	NASH	NASH	S2A4F1	Significant
200	NASH	NASH	S3A3F2	Significant

Table 6: Evaluation of liver Biopsies by the Liver Pathologists in HCV patients without hepatocellular carcinoma

Case	Reference classification	Algorithmic classification	SAF	Disease severity
1	steatosis	steatosis	S1A1F1	Mild
2	steatosis	steatosis	S1A1F1	Mild
3	NASH	NASH	S1A4F3	Significant
4	NASH	NASH	S3A1F2	Significant
5	NASH	NASH	S2A4F1	Significant
6	steatosis	steatosis	S1A1F1	Mild
7	steatosis	steatosis	S1A1F1	Mild
8	steatosis	steatosis	S1A1F1	Mild
9	NASH	NASH	S3A1F2	Significant
10	NASH	NASH	S2A4F1	Significant
11	steatosis	steatosis	S1A0F0	Mild
12	steatosis	steatosis	S1A1F1	Mild
13	steatosis	steatosis	S1A1F1	Mild
14	steatosis	steatosis	S1A1F1	Mild
15	NASH	NASH	S3A1F2	Significant
16	steatosis	steatosis	S1A1F1	Mild
17	steatosis	steatosis	S1A1F1	Mild
18	NASH	NASH	S3A4F3	Significant
19	NASH	NASH	S3A3F2	Significant
20	NASH	NASH	S2A4F1	Significant
21	NASH	NASH	S2A4F1	Significant
22	steatosis	steatosis	S1A1F1	Mild
23	steatosis	steatosis	S1A0F0	Mild
24	steatosis	steatosis	S1A1F1	Mild
25	NASH	NASH	S3A3F2	Significant
26	steatosis	steatosis	S2A1F1	Mild
27	steatosis	steatosis	S1A1F1	Mild
28	NASH	NASH	S2A4F1	Significant
29	steatosis	steatosis	S1A0F0	Mild
30	steatosis	steatosis	S1A1F1	Mild
31	steatosis	steatosis	S1A0F0	Mild
32	NASH	NASH	S3A3F2	Significant
33	NASH	NASH	S3A1F2	Significant
34	NASH	NASH	S3A1F2	Significant
35	NASH	NASH	S2A4F1	Significant
36	NASH	NASH	S3A1F2	Significant
37	steatosis	steatosis	S1A1F1	Mild
38	steatosis	steatosis	S1A0F0	Mild
39	steatosis	steatosis	S1A0F0	Mild
40	NASH	NASH	S3A3F2	Significant
41	steatosis	steatosis	S3A1F0	Mild
42	steatosis	steatosis	S3A1F0	Mild
43	steatosis	steatosis	S1A0F0	Mild
44	NASH	NASH	S3A1F2	Significant
45	steatosis	steatosis	S3A1F0	Mild
46	steatosis	steatosis	S1A1F1	Mild
47	NASH	NASH	S3A3F2	Significant
48	NASH	NASH	S3A3F2	Significant
49	NASH	NASH	S2A4F1	Significant
50	steatosis	steatosis	S3A1F0	Mild
51	steatosis	steatosis	S1A1F1	Mild
52	steatosis	steatosis	S3A1F0	Mild
53	NASH	NASH	S2A4F1	Significant
54	steatosis	steatosis	S1A1F1	Mild
55	NASH	NASH	S3A1F2	Significant
56	NASH	NASH	S2A4F1	Significant
57	steatosis	steatosis	S2A1F1	Mild
58	steatosis	steatosis	S1A1F1	Mild
59	NASH	NASH	S3A1F2	Significant
60	steatosis	steatosis	S2A1F1	Mild
61	steatosis	steatosis	S2A1F1	Mild
62	NASH	NASH	S2A4F1	Significant
63	NASH	NASH	S3A3F2	Significant
64	NASH	NASH	S3A1F2	Significant
65	NASH	NASH	S2A4F1	Significant
66	steatosis	steatosis	S1A1F1	Mild
67	steatosis	steatosis	S2A1F1	Mild
68	steatosis	steatosis	S1A1F1	Mild

*Reference interpretation: is the initial evaluation done by pathologist using accepted criteria, algorithm and SAF score.

*Algorithmic Classification: (Steatosis vs. NASH) after using the algorithm.

* Based on SAF score: mild for A < 2 and F < 2 or significant for A > 2 and/or F > 2.

Table 6: Evaluation of liver Biopsies by the Liver Pathologists in HCV patients without hepatocellular carcinoma

69	NASH	NASH	S2A4F1	Significant
70	steatosis	steatosis	S1A0F0	Mild
71	NASH	NASH	S3A1F2	Significant
72	NASH	NASH	S2A4F1	Significant
73	steatosis	steatosis	S1A0F0	Mild
74	steatosis	steatosis	S1A1F1	Mild
75	steatosis	steatosis	S1A1F1	Mild
76	steatosis	steatosis	S1A1F1	Mild
77	NASH	NASH	S3A1F2	Significant
78	steatosis	steatosis	S1A1F1	Mild
79	steatosis	steatosis	S1A1F1	Mild
80	NASH	NASH	S3A4F3	Significant
81	NASH	NASH	S3A3F2	Significant
82	NASH	NASH	S2A4F1	Significant
83	NASH	NASH	S3A1F2	Significant
84	NASH	NASH	S2A4F1	Significant
85	steatosis	steatosis	S1A0F0	Mild
86	NASH	NASH	S2A4F1	Significant
87	steatosis	steatosis	S1A0F0	Mild
88	NASH	NASH	S3A1F2	Significant
89	NASH	NASH	S2A4F1	Significant
90	steatosis	steatosis	S1A0F0	Mild
91	steatosis	steatosis	S1A1F1	Mild
92	steatosis	steatosis	S1A1F1	Mild
93	steatosis	steatosis	S1A1F1	Mild
94	NASH	NASH	S3A1F2	Significant
95	NASH	NASH	S2A4F1	Significant
96	steatosis	steatosis	S1A0F0	Mild
97	steatosis	steatosis	S1A0F0	Mild
98	NASH	NASH	S3A1F2	Significant
99	NASH	NASH	S2A4F1	Significant
100	steatosis	steatosis	S1A0F0	Mild
101	steatosis	steatosis	S1A1F1	Mild
102	steatosis	steatosis	S1A1F1	Mild
103	steatosis	steatosis	S1A1F1	Mild
104	NASH	NASH	S3A1F2	Significant
105	steatosis	steatosis	S1A1F1	Mild
106	steatosis	steatosis	S1A1F1	Mild
107	NASH	NASH	S3A4F3	Significant
108	NASH	NASH	S3A3F2	Significant
109	NASH	NASH	S2A4F1	Significant
110	NASH	NASH	S3A1F2	Significant
111	NASH	NASH	S2A4F1	Significant
112	steatosis	steatosis	S1A0F0	Mild
113	NASH	NASH	S2A4F1	Significant
114	steatosis	steatosis	S1A0F0	Mild
115	NASH	NASH	S3A1F2	Significant
116	NASH	NASH	S2A4F1	Significant
117	steatosis	steatosis	S1A0F0	Mild
118	steatosis	steatosis	S1A1F1	Mild
119	steatosis	steatosis	S1A1F1	Mild
120	steatosis	steatosis	S1A1F1	Mild
121	NASH	NASH	S3A1F2	Significant
122	NASH	NASH	S2A4F1	Significant
123	steatosis	steatosis	S1A0F0	Mild
124	NASH	NASH	S3A3F2	Significant
125	NASH	NASH	S2A4F1	Significant
126	NASH	NASH	S3A1F2	Significant
127	NASH	NASH	S2A4F1	Significant
128	steatosis	steatosis	S1A0F0	Mild
129	NASH	NASH	S2A4F1	Significant
130	steatosis	steatosis	S1A0F0	Mild
131	NASH	NASH	S3A1F2	Significant
132	NASH	NASH	S2A4F1	Significant
133	steatosis	steatosis	S1A0F0	Mild
134	steatosis	steatosis	S1A1F1	Mild
135	steatosis	steatosis	S1A1F1	Mild
136	steatosis	steatosis	S1A1F1	Mild
137	NASH	NASH	S3A1F2	Significant
138	NASH	NASH	S2A4F1	Significant

*Reference interpretation: is the initial evaluation done by pathologist using accepted criteria, algorithm and SAF score.

*Algorithmic Classification: (Steatosis vs. NASH) after using the algorithm.

* Based on SAF score: mild for A < 2 and F < 2 or significant for A > 2 and/or F > 2.

Table 6: Evaluation of liver Biopsies by the Liver Pathologists in HCV patients without hepatocellular carcinoma

139	steatosis	steatosis	S1A0F0	Mild
140	steatosis	steatosis	S1A0F0	Mild
141	NASH	NASH	S3A1F2	Significant
142	NASH	NASH	S2A4F1	Significant
143	steatosis	steatosis	S1A0F0	Mild
144	steatosis	steatosis	S1A1F1	Mild
145	NASH	NASH	S3A1F2	Significant
146	NASH	NASH	S2A4F1	Significant
147	steatosis	steatosis	S1A0F0	Mild
148	NASH	NASH	S3A3F2	Significant
149	NASH	NASH	S2A4F1	Significant
150	NASH	NASH	S3A1F2	Significant
151	NASH	NASH	S2A4F1	Significant
152	steatosis	steatosis	S1A0F0	Mild
153	NASH	NASH	S2A4F1	Significant
154	steatosis	steatosis	S1A0F0	Mild
155	NASH	NASH	S3A1F2	Significant
156	NASH	NASH	S2A4F1	Significant
157	steatosis	steatosis	S1A0F0	Mild
158	steatosis	steatosis	S1A1F1	Mild
159	steatosis	steatosis	S1A1F1	Mild
160	steatosis	steatosis	S1A1F1	Mild
161	NASH	NASH	S3A1F2	Significant
162	NASH	NASH	S2A4F1	Significant
163	steatosis	steatosis	S1A0F0	Mild
164	steatosis	steatosis	S1A0F0	Mild
165	NASH	NASH	S3A1F2	Significant
166	NASH	NASH	S2A4F1	Significant
167	steatosis	steatosis	S1A0F0	Mild
168	steatosis	steatosis	S1A1F1	Mild
169	steatosis	steatosis	S1A0F0	Mild
170	NASH	NASH	S3A3F2	Significant
171	NASH	NASH	S2A4F1	Significant
172	NASH	NASH	S3A1F2	Significant
173	NASH	NASH	S2A4F1	Significant
174	steatosis	steatosis	S1A0F0	Mild
175	NASH	NASH	S2A4F1	Significant
176	steatosis	steatosis	S1A0F0	Mild
177	NASH	NASH	S3A1F2	Significant
178	NASH	NASH	S2A4F1	Significant
179	steatosis	steatosis	S1A0F0	Mild
180	steatosis	steatosis	S1A1F1	Mild
181	steatosis	steatosis	S1A1F1	Mild
182	steatosis	steatosis	S1A1F1	Mild
183	NASH	NASH	S3A1F2	Significant
184	NASH	NASH	S2A4F1	Significant
185	steatosis	steatosis	S1A0F0	Mild
186	NASH	NASH	S3A3F2	Significant
187	NASH	NASH	S2A4F1	Significant
188	NASH	NASH	S3A1F2	Significant
189	NASH	NASH	S2A4F1	Significant
190	steatosis	steatosis	S1A0F0	Mild
191	NASH	NASH	S2A4F1	Significant
192	steatosis	steatosis	S1A0F0	Mild
193	NASH	NASH	S3A1F2	Significant
194	NASH	NASH	S2A4F1	Significant
195	steatosis	steatosis	S1A0F0	Mild
196	steatosis	steatosis	S1A1F1	Mild
197	steatosis	steatosis	S1A1F1	Mild
198	steatosis	steatosis	S1A1F1	Mild
199	NASH	NASH	S3A1F2	Significant
200	NASH	NASH	S2A4F1	Significant

*Reference interpretation: is the initial evaluation done by pathologist using accepted criteria, algorithm and SAF score.

*Algorithmic Classification: (Steatosis vs. NASH) after using the algorithm.

* Based on SAF score: mild for A < 2 and F < 2 or significant for A > 2 and/or F > 2.

Table 7

Odds ratio (OR) and 95% confidence interval (CI) of clinical status and of CD 24 C/T SNP 170 genotypic frequencies in 400 HCV patients.

Variable	Genotypic frequencies		OR (95% CI)	<i>p</i> value
	CC	CT + TT		
Clinical Stage				
Stage I/II	160 (45.2%)	38 (53.4%)	1.00	<i>p</i> =0.537
Stage III/IV	30 (32.7%)	88 (78.8%)	1.121 (0.371-1.640)	
Lymph node metastasis				
No	230(83.2%)	64(74.4%)	1.00	* <i>p</i> =0.627
Yes	2 (3.1%)	10 (4.6%)	0.722 (0.158-2.568)	
Distant metastasis				
No	220 (85.1%)	65 (83.4%)	1.00	<i>p</i> =0.526
Yes	9 (4.9%)	16 (4.6%)	1.453 (0.542-2.786)	
Vascular invasion				
No	220 (82.5%)	60 (55.9%)	1.00	* <i>p</i> =0.452
Yes	41 (16.5%)	200 (83.1%)	1.521 (0.652-1.987)	
Liver cirrhosis				
Negative	52 (19.8%)	12 (19.5%)	1.00	* <i>p</i> =0.821
Positive	80 (20.1%)	320 (80.2%)	1.562 (0.634-1.823)	

The ORs with analyzed by their 95% CIs were estimated by logistic regression models.

* *p* value < 0.05 as statistically significant.






Fig 1: A photomicrograph in control human liver section showing normal architecture of hepatic lobules, in the form of hepatocytes arranged as plates radiating from the central vein. The liver plates were separated from each other by irregular sinusoids. The hepatocytes appeared polyhedral in shape with acidophilic cytoplasm with large, rounded & vesicular nuclei (Hx&E X100)


Preprint



A)



B)



C)

Fig 2: (A): A photomicrograph in human liver section with hepatitis C infection showing Mallory-Denk bodies (arrow) & fatty infiltration (Hx&E.40x). (B): A photomicrograph in human liver section with hepatitis C infection showing Megamitochondria (arrow head) in a cell with microvesicular steatosis (Hx&E 40x). (C): A photomicrograph of a human liver section with NASH showing Peri-sinusoidal fibrosis (Masson's trichrome, 20x)




Fig 3: A photomicrograph in human liver sections with hepatitis C infection & hepatocellular carcinoma showing: (A): Masses of malignant cells with frequent mitosis, hyper chromatic nuclei & trabecular growth pattern (HX&E 100x). (B): Pseudo glandular growth pattern (HX&E 100X) (C): Loss of architecture, cellular degeneration & Solid growth pattern (HX&E 200x). (D): Loss of architecture with dilated central vein & giant cell formation (HX&E 200X)




Figure 4: Correlation between frequency of CD24 polymorphism and tumor status in HCC patients
P < 0.0001




Figure 5: Kaplan-Meier curves for probability of survival by CD24 polymorphism among groups at $P < 0.0001$

Preprint




Figure 6: Column scatter dot plot showing Talin-1 gene expression among control, HCV, and HCV+HCC groups, $P < 0.0001$

Preprint





Figure 7: Correlation of CD24 170 SNP genotypes with Taline-1 gene expression in HCV patients.
 $P < 0.0001$

Preprint



1- Specificity%
Figure 8: AUC curve analysis of talin-1 gene expression ,P <0.0001, AUC=0.9